



MATERNAL MENTAL HEALTH: PERINATAL DEPRESSION AND ANXIETY

READINESS

Every Clinical Care Setting

- Identify mental health screening tools to be made available in every clinical setting (outpatient OB clinics and inpatient facilities).
- Establish a response protocol and identify screening tools for use based on local resources.
- Educate clinicians and office staff on use of the identified screening tools and response protocol.
- Identify an individual who is responsible for driving adoption of the identified screening tools and response protocol.

RECOGNITION & PREVENTION

Every Woman

- Obtain individual and family mental health history (including past and current medications) at intake, with review and update as needed.
- Conduct validated mental health screening during appropriately timed patient encounters, to include both during pregnancy and in the postpartum period.
- Provide appropriately timed perinatal depression and anxiety awareness education to women and family members or other support persons.

Standardization of health care processes and reduced variation has been shown to improve outcomes and quality of care. The Council on Patient Safety in Women's Health Care disseminates patient safety bundles to help facilitate the standardization process. This bundle reflects emerging clinical, scientific, and patient safety advances as of the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Although the components of a particular bundle may be adapted to local resources, standardization within an institution is strongly encouraged.

The Council on Patient Safety in Women's Health Care is a broad consortium of organizations across the spectrum of women's health for the promotion of safe health care for every woman.



RESPONSE

Every Case

- Initiate a stage-based response protocol for a positive mental health screen.
- Activate an emergency referral protocol for women with suicidal/homicidal ideation or psychosis.
- Provide appropriate and timely support for women, as well as family members and staff, as needed.
- Obtain follow-up from mental health providers on women referred for treatment. This should include the necessary release of information forms.

REPORTING/SYSTEMS LEARNING

Every Clinical Care Setting

- Establish a non-judgmental culture of safety through multidisciplinary mental health rounds.
- Perform a multidisciplinary review of adverse mental health outcomes.
- Establish local standards for recognition and response in order to measure compliance, understand individual performance, and track outcomes.

Standardization of health care processes and reduced variation has been shown to improve outcomes and quality of care. The Council on Patient Safety in Women's Health Care disseminates patient safety bundles to help facilitate the standardization process. This bundle reflects emerging clinical, scientific, and patient safety advances as of the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Although the components of a particular bundle may be adapted to local resources, standardization within an institution is strongly encouraged.

The Council on Patient Safety in Women's Health Care is a broad consortium of organizations across the spectrum of women's health for the promotion of safe health care for every woman.